

Ehrhart polynomials, Hecke series, and affine buildings

Claudia Alfes^{*1}, and Joshua Maglione^{†2}, and Christopher Voll^{‡1}

¹*Fakultät für Mathematik, Universität Bielefeld, Germany*

²*School of Mathematical and Statistical Sciences, University of Galway, Ireland*

Abstract. Given a lattice polytope P and a prime p , we define a function from the set of primitive symplectic p -adic lattices to the rationals that extracts the ℓ th coefficient of the Ehrhart polynomial of P relative to the given lattice. Inspired by work of Gunnells and Rodriguez Villegas in type A, we show that these functions are eigenfunctions of a suitably defined action of the spherical symplectic Hecke algebra. Although they depend significantly on the polytope P , their eigenvalues are independent of P and expressed as polynomials in p . We define local zeta functions that enumerate the values of these Hecke eigenfunctions on the vertices of the affine Bruhat–Tits buildings associated with p -adic symplectic groups. We compute these zeta functions by enumerating p -adic lattices by their elementary divisors and, simultaneously, one Hermite parameter. We report on a general functional equation satisfied by these local zeta functions, confirming a conjecture of Vankov.

Keywords: Ehrhart polynomials, Hecke series, affine buildings, Satake isomorphism, symplectic lattices

1 Introduction

Let P be a fixed full-dimensional lattice polytope in \mathbb{R}^n , i.e. the convex hull of finitely many points $V(P)$ in $\Lambda_0 = \mathbb{Z}^n$. Given a lattice Λ such that $\Lambda_0 \subseteq \Lambda \subseteq \mathbb{Q}^n$, we denote the Ehrhart polynomial of P with respect to Λ by

$$E^\Lambda(P) = \sum_{\ell=0}^n c_\ell^\Lambda(P) T^\ell \in \mathbb{Q}[T]. \quad (1.1)$$

It is of interest to describe the variation of the coefficients $c_\ell^\Lambda(P)$ with Λ as compared to $c_\ell(P) = c_\ell^{\Lambda_0}(P)$; write $E(P)$ for $E^{\Lambda_0}(P)$. For $g \in \mathrm{GL}_n(\mathbb{Q}) \cap \mathrm{Mat}_n(\mathbb{Z})$ we define

$$g \cdot P = \mathrm{conv}\{g \cdot v \mid v \in V(P)\},$$

^{*}alfes@math.uni-bielefeld.de. Alfes was supported by the Daimler and Benz Foundation.

[†]joshua.maglione@universityofgalway.ie. Maglione was supported by the DFG-GRK 2297.

[‡]C.Voll.98@cantab.net. Alfes and Voll are funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) — SFB-TRR 358/1 2023 — 491392403.

which is again a lattice polytope. We write Λ_g for the lattice generated by the rows of $g \in \mathrm{GL}_n(\mathbb{Q})$. Thus, for every $g \in \mathrm{GL}_n(\mathbb{Q}) \cap \mathrm{Mat}_n(\mathbb{Z})$, we have

$$E(g \cdot P) = E^{\Lambda_{g^{-1}}}(P). \quad (1.2)$$

We note that $\Lambda_g \subseteq \mathbb{Z}^n \subseteq \Lambda_{g^{-1}} \subseteq \mathbb{Q}^n$ for $g \in \mathrm{GL}_n(\mathbb{Q}) \cap \mathrm{Mat}_n(\mathbb{Z})$ with $|\det(g)| > 1$.

Gunnells and Rodriguez Villegas [3] consider how the coefficients of $E^\Lambda(P)$ from Equation (1.1) relate to $E(P)$ for lattices Λ such that $\Lambda_0 \subseteq \Lambda \subseteq p^{-1}\Lambda_0 \subseteq \mathbb{Q}^n$. In Section 2.1 we revisit these results from our perspective. In addition, we consider a symplectic analogue of the work of Gunnells and Rodriguez Villegas.

1.1 Zeta functions of Ehrhart coefficients

For a prime p , we write \mathbb{Z}_p for the ring of p -adic integers and \mathbb{Q}_p for its field of fractions. Below we define, for each $n \in \mathbb{N} = \{1, 2, \dots\}$ and $\ell \in [2n]_0 = \{0, \dots, 2n\}$, local zeta functions which we call *Ehrhart–Hecke zeta functions*. These functions are Dirichlet series in a complex variable s encoding the ratio of ℓ th coefficients of the Ehrhart polynomial of P , as the lattice Λ varies among symplectic lattices in \mathbb{Q}_p^{2n} .

Recall the group scheme GSp_{2n} of symplectic similitudes. For a ring K its K -rational points are, with $J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$,

$$\mathrm{GSp}_{2n}(K) = \{A \in \mathrm{GL}_{2n}(K) \mid AJA^t = \mu(A)J, \text{ for some } \mu(A) \in K^\times\}.$$

We set $G_n = \mathrm{GSp}_{2n}(\mathbb{Q}_p)$, $\Gamma_n = \mathrm{GSp}_{2n}(\mathbb{Z}_p)$, and $G_n^+ = \mathrm{GSp}_{2n}(\mathbb{Q}_p) \cap \mathrm{Mat}_{2n}(\mathbb{Z}_p)$. The set G_n^+/Γ_n is in bijection with the set of special vertices of the affine building associated with the group $\mathrm{GSp}_{2n}(\mathbb{Q}_p)$, which is of type \tilde{C}_n .

We define the (*local*) *Ehrhart–Hecke zeta function* (of type C) as

$$\mathcal{Z}_{n,\ell,p}^C(s) = \sum_{g \in G_n^+/\Gamma_n} \frac{c_\ell^{\Lambda_{g^{-1}}}(P)}{c_\ell(P)} |\Lambda_{g^{-1}} : \mathbb{Z}_p^n|^{-s}.$$

Informally speaking, the zeta function $\mathcal{Z}_{n,\ell,p}^C(s)$ hence encodes the average ℓ th coefficient of the Ehrhart polynomial of P across certain symplectic lattices.

1.2 Symplectic Hecke series

The zeta functions of Section 1.1 are closely connected to formal power series over the Hecke algebra associated with the pair (G_n^+, Γ_n) . To explain this connection, we establish additional notation. For $m \in \mathbb{N}$ we define

$$D_n^C(m) = \{A \in G_n^+ \mid AJA^t = mJ\}.$$

Let $\mathcal{H}_p^C = \mathcal{H}^C(G_n^+, \Gamma_n)$ be the spherical Hecke algebra. The Hecke operator $T_n^C(m)$ is

$$T_n^C(m) = \sum_{g \in \Gamma_n \backslash D_n^C(m) / \Gamma_n} \Gamma_n g \Gamma_n.$$

The (formal) symplectic Hecke series is defined as

$$\sum_{\alpha \geq 0} T_n^C(p^\alpha) X^\alpha \in \mathcal{H}_p^C[[X]]. \quad (1.3)$$

Shimura's conjecture [6] that the series in (1.3) is a rational function in X was proved by Andrianov [1]. Explicit formulae, however, seem only to be known for $n \leq 4$; see [9].

We consider the image of the Hecke series in (1.3) under the Satake isomorphism $\Omega : \mathcal{H}_p^C \rightarrow \mathbb{C}[x_0^{\pm 1}, \dots, x_n^{\pm 1}]^W$ mapping onto the ring of invariants of W , the Weyl group of G_n . For variables $\mathbf{x} = (x_0, \dots, x_n)$, we define the (local) Satake generating function as

$$R_{n,p}(\mathbf{x}, X) = \sum_{\alpha \geq 0} \Omega(T_n^C(p^\alpha)) X^\alpha \in \mathbb{C}[\mathbf{x}^{\pm 1}][[X]].$$

and the (local) primitive local Satake generating function as

$$R_{n,p}^{\text{pr}}(\mathbf{x}, X) = (1 - x_0 X) (1 - x_0 x_1 \cdots x_n X) R_{n,p}(\mathbf{x}, X). \quad (1.4)$$

We write $V(\mathcal{X}_n)$ for the set of vertices of \mathcal{X}_n , the affine building \mathcal{X}_n of type \tilde{A}_{n-1} associated with the group $\text{GL}_n(\mathbb{Q}_p)$, viz. homothety classes of full lattices in $\text{GL}_n(\mathbb{Q}_p)$. In [2, Section 3.3] Andrianov shows, in essence, that $R_{n,p}^{\text{pr}}$ can be interpreted as a sum over $V(\mathcal{X})$; see Theorem 1.1 below.

For a lattice $\Lambda \leq \mathbb{Z}_p^n$, set $\mathbf{v}(\Lambda) = (v_1 \leq \dots \leq v_n) \in \mathbb{N}_0^n$ if $\mathbb{Z}_p^n / \Lambda \cong \mathbb{Z} / p^{v_1} \oplus \dots \oplus \mathbb{Z} / p^{v_n}$. Setting $v_0 = 0$, we define

$$\boldsymbol{\mu}(\Lambda) = (\mu_1, \dots, \mu_n) = (v_n - v_{n-1}, \dots, v_1 - v_0).$$

Having chosen a \mathbb{Z}_p -basis of \mathbb{Z}_p^n we associate to each lattice $\Lambda \leq \mathbb{Z}_p^n$ a unique matrix

$$M_\Lambda = \begin{pmatrix} p^{\delta_1} & m_{12} & \cdots & m_{1n} \\ & p^{\delta_2} & \cdots & m_{2n} \\ & & \ddots & \vdots \\ & & & p^{\delta_n} \end{pmatrix} \in \text{Mat}_n(\mathbb{Z}_p), \quad (1.5)$$

whose rows generate Λ and with $0 \leq v_p(m_{ij}) \leq \delta_j$ for all $1 \leq i < j \leq n$. The matrix M_Λ in (1.5) is said to be in Hermite normal form. We set $\delta(\Lambda) = (\delta_1, \dots, \delta_n)$. Clearly each homothety class $[\Lambda]$ contains a unique representative $\Lambda_m \leq \mathbb{Z}_p^n$ such that $p^{-1}\Lambda_m \not\leq \mathbb{Z}_p^n$.

Theorem 1.1 (Andrianov). *Let $n \in \mathbb{N}$, $\mathbf{a} = (1, 2, \dots, n) \in \mathbb{N}^n$, $\mathbf{d} = (n, n-1, \dots, 1) \in \mathbb{N}^n$, and let \langle, \rangle be the usual dot product. Then*

$$R_{n,p}^{\text{pr}}(\mathbf{x}, X) = \sum_{[\Lambda] \in V(\mathcal{X}_n)} p^{\langle \mathbf{d}, \mathbf{v}(\Lambda_m) \rangle - \langle \mathbf{a}, \delta(\Lambda_m) \rangle} x_1^{\delta_1(\Lambda_m)} \cdots x_n^{\delta_n(\Lambda_m)} (x_0 X)^{v_n(\Lambda_m)}.$$

1.3 The Hermite–Smith generating function

We define a generating function enumerating finite-index sublattices of \mathbb{Z}_p^n simultaneously by their Hermite and Smith normal forms. For $n \in \mathbb{N}$, let $\mathbf{X} = (X_1, \dots, X_n)$ and $\mathbf{Y} = (Y_1, \dots, Y_n)$ be variables. The *Hermite–Smith generating function* is

$$\mathrm{HS}_{n,p}(\mathbf{X}, \mathbf{Y}) = \sum_{\Lambda \leq \mathbb{Z}_p^n} \mathbf{X}^{\mu(\Lambda)} \mathbf{Y}^{\delta(\Lambda)} = \sum_{\Lambda \leq \mathbb{Z}_p^n} \prod_{i=1}^n X_i^{\mu_i(\Lambda)} Y_i^{\delta_i(\Lambda)} \in \mathbb{Z}[[\mathbf{X}, \mathbf{Y}]]. \quad (1.6)$$

Clearly, if $\Lambda \leq \mathbb{Z}_p^n$ has finite index, then so does $p^m \Lambda$ for all $m \in \mathbb{N}_0$. This allows us to extract a “homothety factor” from the sum defining $\mathrm{HS}_{n,p}(\mathbf{X}, \mathbf{Y})$. The *primitive Hermite–Smith generating function* is

$$\mathrm{HS}_{n,p}^{\mathrm{pr}}(\mathbf{X}, \mathbf{Y}) = \sum_{[\Lambda] \in \mathcal{V}(\mathcal{X}_n)} \mathbf{X}^{\mu(\Lambda_m)} \mathbf{Y}^{\delta(\Lambda_m)} = (1 - X_n Y_1 \cdots Y_n) \mathrm{HS}_{n,p}(\mathbf{X}, \mathbf{Y}). \quad (1.7)$$

With this generating function we may obtain the primitive local Satake generating function of Section 1.2, as follows. We define a ring homomorphism

$$\begin{aligned} \Phi : \mathbb{Q}[[X_1, X_2, \dots, Y_1, Y_2, \dots]] &\longrightarrow \mathbb{Q}[[x_0, x_1, \dots, X]] \\ X_i &\longmapsto p^{\binom{i+1}{2}} x_0 X, \\ Y_i &\longmapsto p^{-i} x_i \end{aligned} \quad (1.8)$$

for all $i \in \mathbb{N}_0$. By design of Φ and Theorem 1.1 we have $\Phi(\mathrm{HS}_{n,p}^{\mathrm{pr}}) = R_{n,p}^{\mathrm{pr}}$.

Example 1.2. For $n = 2$, the Hermite–Smith generating function is

$$\begin{aligned} \mathrm{HS}_{2,p}(\mathbf{X}, \mathbf{Y}) &= \frac{1 - X_1^2 Y_1 Y_2}{(1 - X_1 Y_1)(1 - p X_1 Y_2)(1 - X_2 Y_1 Y_2)}, \\ R_{2,p}(x, X) &= \frac{1 - p^{-1} x_0^2 x_1 x_2 X^2}{(1 - x_0 X)(1 - x_0 x_1 X)(1 - x_0 x_2 X)(1 - x_0 x_1 x_2 X)}. \end{aligned}$$

2 Main results

Interpreting the ℓ -th coefficients of the Ehrhart polynomial of the polytope P as a function on a set of (homothety classes of) p -adic lattices invites the definition of an action of the spherical Hecke algebra $\mathcal{H}_p^{\mathrm{C}}$. The latter is generated by a set of $n + 1$ generators $T_n^{\mathrm{C}}(p, 0), T_n^{\mathrm{C}}(p^2, 1), \dots, T_n^{\mathrm{C}}(p^2, n)$. It suffices to explain how these generators act. For $k \in [n]$, define diagonal matrices in G_n^+ as follows:

$$D_0 = \mathrm{diag}(\underbrace{1, \dots, 1}_n, \underbrace{p, \dots, p}_n), \quad D_k = \mathrm{diag}(\underbrace{1, \dots, 1}_{n-k}, \underbrace{p, \dots, p}_k, \underbrace{p^2, \dots, p^2}_{n-k}, \underbrace{p, \dots, p}_k).$$

Set $\mathcal{D}_{n,k}^{\mathbb{C}} = \Gamma_n D_k \Gamma_n / \Gamma_n$. The set $\mathcal{D}_{n,k}^{\mathbb{C}}$ can be interpreted as the set of symplectic lattices with symplectic elementary divisors equal to those of D_k . We define

$$T_n^{\mathbb{C}}(p,0)E(P) = \sum_{g \in \mathcal{D}_{n,0}^{\mathbb{C}}} E(g \cdot P), \quad T_n^{\mathbb{C}}(p^2,k)E(P) = \sum_{g \in \mathcal{D}_{n,k}^{\mathbb{C}}} E(g \cdot P).$$

For $\ell \geq \mathbb{N}_0$, we define functions

$$\mathcal{E}_{n,p,\ell,P} : G_n^+ / \Gamma_n \rightarrow \mathbb{C}, \quad \Gamma_n g \mapsto c_\ell(E^{\wedge_{s-1}}(P)).$$

Lastly, for all $T \in \mathcal{H}_p^{\mathbb{C}}$ set

$$T \mathcal{E}_{n,p,\ell,P}(\Gamma_n g) = c_\ell(T E^{\wedge_{s-1}}(P)).$$

Recall that P is full-dimensional; for $k \in [n]$, and $\ell \in [2n]_0$, we define

$$v_{n,0,\ell}^{\mathbb{C}}(p) = \frac{c_\ell(T_n^{\mathbb{C}}(p,0)E(P))}{c_\ell(E(P))}, \quad v_{n,k,\ell}^{\mathbb{C}}(p) = \frac{c_\ell(T_n^{\mathbb{C}}(p^2,k)E(P))}{c_\ell(E(P))}.$$

The notation suggests that the value $v_{n,k,\ell}^{\mathbb{C}}(p)$ is independent of the polytope P , which is justified by [Theorem A](#). General properties of the Ehrhart polynomial imply that

$$v_{n,n,\ell}^{\mathbb{C}}(p) = p^\ell, \quad v_{n,k,0}^{\mathbb{C}}(p) = \#\mathcal{D}_{n,k}^{\mathbb{C}}.$$

Every \mathbb{Q} -linear homomorphism $\lambda : \mathcal{H}_p^{\mathbb{C}} \rightarrow \mathbb{C}$ is uniquely determined by parameters $(a_0, \dots, a_n) \in \mathbb{C}^{n+1}$ such that if $\psi : \mathbb{C}[x_0^{\pm 1}, \dots, x_n^{\pm 1}] \rightarrow \mathbb{C}$ is given by $x_i = a_i$ then $\lambda = \psi \circ \Omega$; see [2, Proposition 3.3.36].

Theorem A. *The functions $\mathcal{E}_{n,p,\ell,P}$ are Hecke eigenfunctions under the action defined above; specifically, for all $k \in [n]$, we have*

$$T_n^{\mathbb{C}}(p,0)\mathcal{E}_{n,p,\ell,P} = v_{n,0,\ell}^{\mathbb{C}}(p)\mathcal{E}_{n,p,\ell,P}, \quad T_n^{\mathbb{C}}(p^2,k)\mathcal{E}_{n,p,\ell,P} = v_{n,k,\ell}^{\mathbb{C}}(p)\mathcal{E}_{n,p,\ell,P},$$

where the $v_{n,k,\ell}^{\mathbb{C}}(p)$ are polynomials in p with integer coefficients which are independent of P . Moreover, the parameters associated to $v_{n,k,\ell}^{\mathbb{C}}(p)$ are $(p^\ell, p, p^2, \dots, p^{n-1}, p^{n-\ell})$.

[Table 1](#) lists the values of $v_{n,k,\ell}^{\mathbb{C}}(p)$ for small values of n and k .

[Theorem A](#) enables us to relate $\mathcal{Z}_{n,\ell,p}^{\mathbb{C}}(s)$ to $R_{n,p}(\mathbf{x}, X)$. Let $\psi_{n,\ell}$ be the ring homomorphism from $\mathbb{C}[[x_0, x_1, \dots, X]] \rightarrow \mathbb{C}[t]$ given by

$$X \mapsto t^n, \quad x_0 \mapsto p^\ell, \quad x_n \mapsto p^{n-\ell}, \quad x_i \mapsto p^i.$$

Corollary B. *For $n \in \mathbb{N}$ and $\ell \in [2n]_0$ we have, writing $t = p^{-s}$,*

$$(\psi_{n,\ell} \circ \Phi)(\text{HS}_{n,p}^{\text{Pr}}(\mathbf{X}, \mathbf{Y})) = \psi_{n,\ell}(R_{n,p}^{\text{Pr}}) = \mathcal{Z}_{n,\ell,p}^{\mathbb{C}}(s) \left(1 - p^{\ell-s}\right) \left(1 - p^{\binom{n+1}{2}-s}\right).$$

ℓ	$v_{2,0,\ell}^{\mathbb{C}}(p)$	$v_{2,1,\ell}^{\mathbb{C}}(p)$
4	$p^5 + p^4 + p^3 + p^2$	$p^8 + p^7 + p^6 + p^5$
3	$p^4 + p^3 + p^3 + p^2$	$2p^6 + p^5 + 2p^4 - p^3$
2	$p^3 + p^3 + p^2 + p^2$	$p^5 + 3p^4 + p^3 - p^2$
1	$p^3 + p^2 + p^2 + p^1$	$2p^4 + p^3 + 2p^2 - p$
0	$p^3 + p^2 + p + 1$	$p^4 + p^3 + p^2 + p$

Table 1: The polynomials $v_{2,k,\ell}^{\mathbb{C}}(p)$ for $k \in \{0,1\}$ and $\ell \in [4]_0$.

Thanks to [Corollary B](#), we can work with $\text{HS}_{n,p}$ to prove that $R_{n,p}$ and $\mathcal{Z}_{n,\ell,p}^{\mathbb{C}}$ satisfy a self-reciprocity property, which proves the conjecture in [9, Remark 4].

Theorem C. *Let $n \in \mathbb{N}$. Then $\text{HS}_{n,p}(\mathbf{X}, \mathbf{Y})$ is a rational function in \mathbf{X} and \mathbf{Y} . Furthermore, for $\mathbf{X}^{-1} = (X_1^{-1}, \dots, X_n^{-1})$ and $\mathbf{Y}^{-1} = (Y_1^{-1}, \dots, Y_n^{-1})$, we have*

$$\text{HS}_{n,p}(\mathbf{X}^{-1}, \mathbf{Y}^{-1}) \Big|_{p \rightarrow p^{-1}} = (-1)^n p^{\binom{n}{2}} X_n Y_1 \cdots Y_n \cdot \text{HS}_{n,p}(\mathbf{X}, \mathbf{Y}).$$

We prove [Theorem C](#) by writing $\text{HS}_{n,p}$ as a p -adic integral and applying results of [10], where the operation of inverting p is also explained.

Corollary D. *For $n \in \mathbb{N}$ and $\ell \in [2n]_0$, we have*

$$\begin{aligned} \mathcal{Z}_{n,\ell,p}^{\mathbb{C}}(s) \Big|_{p \rightarrow p^{-1}} &= (-1)^{n+1} p^{n^2 + \ell - 2ns} \cdot \mathcal{Z}_{n,\ell,p}^{\mathbb{C}}(s), \\ R_{n,p}(\mathbf{x}, X) \Big|_{p \rightarrow p^{-1}} &= (-1)^{n+1} p^{\binom{n}{2}} x_0^2 x_1 \cdots x_n X^2 \cdot R_{n,p}(\mathbf{x}, X). \end{aligned}$$

In the next theorem, we determine a formula for the specialization of $\text{HS}_{n,p}^{\text{PR}}$ which yields $\mathcal{Z}_{n,\ell,p}^{\mathbb{C}}$ by [Corollary B](#). To this end we define

$$\overline{\text{HS}}_{n,p}(\mathbf{X}, \mathbf{Y}) = \text{HS}_{n,p}^{\text{PR}}(\mathbf{X}, 1, \dots, 1, \mathbf{Y}).$$

We prove that $\overline{\text{HS}}_{n,p}$ is a rational function in the $n+1$ variables \mathbf{X} and \mathbf{Y} and, in addition, the prime p . In order to describe the formula, we define additional notation. For $I = \{i_1 < \cdots < i_\ell\} \subseteq [n-1]$, with $i_{\ell+1} = n$, $k \in [\ell+1]$, and a variable Z , we set

$$\begin{aligned} I^{(k)} &= \{i_j \mid j < k\} \cup \{i_j - 1 \mid j \geq k\} \\ \mathcal{G}_{n,I,k}(Z, \mathbf{X}, \mathbf{Y}) &= \left(\prod_{j=1}^{k-1} \frac{Z^{i_j(n-i_j-1)} X_{i_j}}{1 - Z^{i_j(n-i_j-1)} X_{i_j}} \right) \left(\prod_{j=k}^{\ell} \frac{Z^{i_j(n-i_j)} X_{i_j} Y}{1 - Z^{i_j(n-i_j)} X_{i_j} Y} \right). \end{aligned}$$

Theorem E. Let $n \in \mathbb{N}$. For $I = \{i_1 < \cdots < i_\ell\} \subset [n-1]$, set

$$W_{n,I}(Z, \mathbf{X}, Y) = \sum_{k=1}^{\ell+1} Z^{-(n-i_k)} \binom{n-1}{I^{(k)}}_{Z^{-1}} \mathcal{G}_{n,I,k}(Z, \mathbf{X}, Y) \\ + \sum_{k=1}^{\ell} \frac{(1 - Z^{-i_j}) \mathcal{G}_{n,I,k}(Z, \mathbf{X}, Y)}{1 - Z^{i_j(n-i_j-1)} X_{i_j}} \left(\sum_{m=k+1}^{\ell+1} Z^{-(n-i_m)} \right) \binom{n-1}{I^{(k+1)}}_{Z^{-1}}.$$

Then

$$\overline{\text{HS}}_{n,p}(\mathbf{X}, Y) = \sum_{I \subseteq [n-1]} W_{n,I}(p, \mathbf{X}, Y) \in \mathbb{Z}(p, \mathbf{X}, Y).$$

Via the various substitutions given above, [Theorem E](#) yields explicit formulae for the functions $R_{n,p}$ and, specifically,

$$Z_{n,\ell,p}^{\text{C}}(s) = (1 - p^{\ell-s})^{-1} (1 - p^{\binom{n+1}{2}-s})^{-1} \sum_{I \subseteq [n-1]} W_{n,I} \left(p, \left(p^{\binom{i+1}{2} + \ell - ns} \right)_{i=1}^n, p^{-\ell} \right).$$

In the next theorem we show that the primitive local Satake generating function can be viewed as a “ p -analogue” of the fine Hilbert series of a Stanley–Reisner ring. Let V be a finite set. If $\Delta \subseteq 2^V$ is a simplicial complex on V , then the Stanley–Reisner ring of Δ over a ring K is

$$K[\Delta] = K[X_v \mid v \in V] / \left(\prod_{v \in \sigma} X_v \mid \sigma \in 2^V \setminus \Delta \right).$$

Theorem F. For all $n \in \mathbb{N}$, let Δ_n be the n -simplex with vertices $[n]$ and $\Delta = \text{sd}(\partial\Delta_n)$, the barycentric subdivision of boundary of Δ_n , with vertices given by the nonempty subsets of $[n]$. Let $\mathbf{y} = (y_I : \emptyset \neq I \subseteq [n])$ and $\varphi : \mathbb{Z}[\mathbf{y}] \rightarrow \mathbb{Z}[\mathbf{x}, X]$ via $y_I \mapsto x_0 X \prod_{i \in I} x_i$. Then

$$R_{n,p}^{\text{pr}}(\mathbf{x}, X) \Big|_{p \rightarrow 1} = \varphi(\text{Hilb}(\mathbb{Z}[\Delta]; \mathbf{y})) = \sum_{\sigma \in \Delta} \prod_{J \in \sigma} \frac{\varphi(y_J)}{1 - \varphi(y_J)}.$$

With [Theorem F](#), we come full circle and relate the local Satake generating function $R_{n,p}$ to the Ehrhart series of the n -cube.

Corollary 2.1. For all $n \in \mathbb{N}$, let P be the n -cube. Then

$$R_{n,p}(\mathbf{1}, X) \Big|_{p \rightarrow 1} = \text{Ehr}_P(X) = \frac{E_n(X)}{(1 - X)^{n+1}},$$

where $E_n(X) = \sum_{\sigma \in S_n} X^{\text{des}(\sigma)}$ is the Eulerian polynomial.

Proof. It follows from [Theorem F](#) that

$$(1 - X)^2 R_{n,p}(\mathbf{1}, X) \Big|_{p \rightarrow 1} = \sum_{\sigma \in \Delta} \prod_{J \in \sigma} \frac{X}{1 - X}, \quad (2.1)$$

where Δ is the barycentric subdivision of the boundary of the n -simplex. From [5, Theore. 9.1] and Equation (2.1) it follows that

$$R_{n,p}(\mathbf{1}, X)|_{p \rightarrow 1} = \frac{E_n(X)}{(1-X)^{n+1}} = \sum_{k \geq 0} (k+1)^n X^k = \text{Ehr}_P(X). \quad \square$$

2.1 The type-A story

Our work was inspired by Gunnells and Rodriguez Villegas. In [3] they considered type-A versions of some of the questions outlined above. We paraphrase parts of [3] from the perspective of our work in type C. For a prime p we define the (*local*) *Ehrhart–Hecke zeta function* (of type A) as

$$\mathcal{Z}_{n,\ell,p}^A(s) = \sum_{\substack{\mathbb{Z}_p^n \leq \Lambda \leq \mathbb{Q}_p^n \\ |\Lambda : \mathbb{Z}_p^n| < \infty}} \frac{c_\ell^\Lambda(P)}{c_\ell(P)} |\Lambda : \mathbb{Z}_p^n|^{-s}. \quad (2.2)$$

Let $\Gamma_n^A = \text{GL}_n(\mathbb{Z})$ and $G_n^A = \text{Mat}_n(\mathbb{Z}) \cap \text{GL}_n(\mathbb{Q})$. For $m \in \mathbb{N}$, let

$$D_n^A(m) = \{g \in G_n^A \mid |\det(g)| = m\},$$

so $D_n^A(m)$ is a finite union of double cosets relative to Γ_n^A . We define

$$T_n^A(m) = \sum_{g \in \Gamma_n^A \backslash D_n^A(m) / \Gamma_n^A} \Gamma_n^A g \Gamma_n^A,$$

where the sum runs over a set of representatives of the double cosets, which is an element of the Hecke algebra determined by (Γ_n^A, G_n^A) . Moreover, if $\gcd(m, m') = 1$, then

$$T_n^A(m) T_n^A(m') = T_n^A(mm').$$

For $k \in [n]_0$ define $\pi_k(p) = \text{diag}(1, \dots, 1, \overbrace{p, \dots, p}^k)$ and $T_n^A(p, k) = \Gamma_n^A \pi_k(p) \Gamma_n^A$, which decomposes into a finite (disjoint) union of right cosets relative to Γ_n^A .

Gunnells and Rodriguez Villegas [3] considered the following action of the Hecke algebra on the Ehrhart polynomial $E(P) = E^{\Lambda_0}(P)$ of P :

$$T_n^A(p, k) E(P) = \sum_{g \in \Gamma_n^A \pi_k(p) \Gamma_n^A / \Gamma_n^A} E(g \cdot P), \quad (2.3)$$

where the sum runs over a set of right coset representatives. The action in (2.3) is independent of the chosen representatives since Γ_n^A comprises bijections of \mathbb{Z}^n . Our definition in (2.3) differs from [3] only cosmetically via (1.2).

Denote by $\text{Gr}(\ell, n, p)$ the set of ℓ -dimensional subspaces in \mathbb{F}_p^n . For $n \in \mathbb{N}$, $\ell, k \in [n]_0$, and $U \in \text{Gr}(\ell, n, p)$, define

$$v_{n,k,\ell}^A(p) = \sum_{W \in \text{Gr}(k,n,p)} \#(U \cap W).$$

Let $\psi_{n,\ell}^A : \mathbb{Q}[x_1^{\pm 1}, \dots, x_n^{\pm 1}] \rightarrow \mathbb{Q}$ be given by $x_n \mapsto p^\ell$ and $x_i \mapsto p^i$ for all $i \in [n-1]$. Let further ω denote the Satake isomorphism from the p -primary part of the Hecke algebra associated with (Γ_n^A, G_n^A) , written \mathcal{H}_p^A , to the symmetric subring of $\mathbb{Q}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$.

Let $s_{n,k}(x_1, \dots, x_n)$ be the (homogeneous) elementary symmetric polynomial of degree k , and set $s_{n,-1} = 0$.

Theorem 2.2 ([3]). *For $n \in \mathbb{N}$, $k, \ell \in [n]_0$, and a prime p , we have*

$$v_{n,k,\ell}^A(p) = p^k \binom{n-1}{k}_p + p^\ell \binom{n-1}{k-1}_p = \psi_{n,\ell}^A(\omega(T_n^A(p, k))).$$

Moreover,

$$z_{n,\ell,p}^A(s) = (1 - p^{\ell-s})^{-1} \prod_{k=1}^{n-1} (1 - p^{k-s})^{-1}.$$

Proof. First we prove the claims concerning $v_{n,k,\ell}^A(p)$. Therefore,

$$\begin{aligned} v_{n,k,\ell}^A(p) &= \binom{n}{k}_p - \binom{n-1}{k-1}_p + p^\ell \binom{n-1}{k-1}_p && \text{([3, Lem. 3.3])} \\ &= p^k \binom{n-1}{k}_p + p^\ell \binom{n-1}{k-1}_p && \text{(Pascal identity)} \\ &= p^k s_{n-1,k}(1, p, \dots, p^{n-2}) + p^\ell s_{n-1,k-1}(1, p, \dots, p^{n-2}) && \text{([4, Ex. I.2.3])} \\ &= p^{-\binom{k}{2}} \psi_{n,\ell}^A(s_{n,k}) \\ &= \psi_{n,\ell}^A(\omega(T_n^A(p, k))). && \text{([2, Lem. 3.2.21])} \end{aligned}$$

We now tend to the last claim. Tamagawa [7] established the identity

$$\sum_{m \geq 0} T_n^A(p^m) X^m = \left(\sum_{k=0}^n (-1)^k p^{\binom{k}{2}} T_n^A(p, k) X^k \right)^{-1} \in \mathcal{H}_p^A[[X]]. \quad (2.4)$$

Applying $\psi_{n,\ell}^A \circ \omega$ to (2.4) and setting $X = p^{-s}$, we have

$$\sum_{m \geq 0} \psi_{n,\ell}^A(\omega(T_n^A(p^m))) p^{-ms} = \left(\sum_{k=0}^n \psi_{n,\ell}^A(s_{n,k}) (-p)^{-ks} \right)^{-1} = (1 - p^{\ell-s})^{-1} \prod_{k=1}^{n-1} (1 - p^{k-s})^{-1}.$$

Since $v_{n,k,\ell}^A(p)$ is an eigenvalue for $T_n(p, k)$, it follows that

$$z_{n,\ell,p}^A(s) = \sum_{m \geq 0} \psi_{n,\ell}^A(\omega(T_n^A(p^m))) p^{-ms}. \quad \square$$

Corollary 2.3. *Let $\zeta(s)$ be the Riemann zeta function. For $n \in \mathbb{N}$ and $\ell \in [n]_0$, we have*

$$\prod_{\text{prime } p} \mathcal{Z}_{n,\ell,p}^A(s) = \zeta(s - \ell) \prod_{k=1}^{n-1} \zeta(s - k).$$

3 Examples

3.1 Hecke eigenfunctions

We give some explicit examples, showing in [Figure 3.1](#) that the eigenfunctions of [Theorem A](#) depend significantly on the polytope. We do this by displaying a graph whose vertices correspond to homothety classes of lattices. We evaluate the functions $\mathcal{E}_{n,p,\ell}$ on Λ_m for each homothety class $[\Lambda]$.

3.2 Local Ehrhart–Hecke zeta functions

For $n \in [3]$ and $\ell \in [2n]_0$, we record the rational functions $W_{n,\ell}(X, Y) \in \mathbb{Q}(X, Y)$ where, for all primes, $\mathcal{Z}_{n,\ell,p}^C(s) = W_{n,\ell}(p, p^{-ns})$. We computed these with SageMath [\[8\]](#).

$$\begin{aligned} W_{1,\ell}(X, Y) &= \frac{1}{(1 - XY)(1 - X^\ell Y)} \\ W_{2,\ell}(X, Y) &= \frac{1 - X^{2+\ell} Y^2}{(1 - X^2 Y)(1 - X^3 Y)(1 - X^\ell Y)(1 - X^{\ell+1} Y)} \\ W_{3,\ell}(X, Y) &= \frac{1 + (X^{1+\ell} + X^4)Y - A_\ell(X)Y^2 + (X^{6+2\ell} + X^{9+\ell})Y^3 + X^{10+2\ell}Y^4}{(1 - X^3 Y)(1 - X^5 Y)(1 - X^6 Y)(1 - X^\ell Y)(1 - X^{2+\ell} Y)(1 - X^{3+\ell} Y)} \\ W_{4,\ell}(X, Y) &= \frac{N_{4,\ell}(X, Y)}{D_{4,\ell}(X, Y)}, \end{aligned}$$

where $A_\ell(X) = X^{7+\ell} + 2X^{6+\ell} + 2X^{4+\ell} + X^{3+\ell}$,

$$\begin{aligned} N_{4,\ell}(X, Y) &= 1 + (X^5 + X^6 + X^7 + X^8 + X^{1+\ell} + X^{2+\ell} + X^{3+\ell} + X^{4+\ell})Y + (X^{13} \\ &\quad - X^{4+\ell} - 2X^{5+\ell} - 2X^{6+\ell} - 2X^{7+\ell} - 2X^{8+\ell} - 2X^{9+\ell} - 3X^{10+\ell} \\ &\quad - 2X^{11+\ell} - 2X^{12+\ell} - 2X^{13+\ell} - X^{14+\ell} + X^{5+2\ell})Y^2 + (X^{14+\ell} \\ &\quad - X^{18+\ell} + X^{10+2\ell} - X^{14+2\ell})Y^3 - (X^{23+\ell} - X^{14+2\ell} - 2X^{15+2\ell} \\ &\quad - 2X^{16+2\ell} - 2X^{17+2\ell} - 3X^{18+2\ell} - 2X^{19+2\ell} - 2X^{20+2\ell} - 2X^{21+2\ell} \\ &\quad - 2X^{22+2\ell} - 2X^{23+2\ell} - X^{24+2\ell} + X^{15+3\ell})Y^4 - (X^{24+2\ell} + X^{25+2\ell} \\ &\quad + X^{26+2\ell} + X^{27+2\ell} + X^{20+3\ell} + X^{21+3\ell} + X^{22+3\ell} + X^{23+3\ell})Y^5 \\ &\quad - X^{28+3\ell}Y^6, \end{aligned}$$

$$D_{4,\ell}(X, Y) = (1 - X^4Y)(1 - X^7Y)(1 - X^9Y)(1 - X^{10}Y) \\ \times (1 - X^\ell Y)(1 - X^{3+\ell}Y)(1 - X^{5+\ell}Y)(1 - X^{6+\ell}Y).$$




Figure 3.1: Polytopes and some values of $\mathcal{E}_{2,2,1,P}$ displayed on lattices in the affine building of type \tilde{A}_1 associated with the group $\mathrm{GSp}_2(\mathbb{Q}_p) \cong \mathrm{GL}_2(\mathbb{Q}_p)$. The center vertex corresponds to the homothety class of the identity, and the values are the linear coefficients of the Ehrhart polynomials with respect to the corresponding lattices.

References

- [1] A. N. Andrianov. “Rationality theorems for Hecke series and Zeta functions of the groups GL_n and Sp_n over local fields”. *Izv. Akad. Nauk SSSR Ser. Mat.* (1969), pp. 466–505.
- [2] A. N. Andrianov. *Quadratic forms and Hecke operators*. Vol. 286. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1987, pp. xii+374. [DOI](#).
- [3] P. E. Gunnells and F. Rodriguez Villegas. “Lattice polytopes, Hecke operators, and the Ehrhart polynomial”. *Selecta Math. (N.S.)* **13.2** (2007), pp. 253–276.
- [4] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Second. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1995, pp. x+475.
- [5] T. K. Petersen. *Eulerian numbers*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser/Springer, New York, 2015, pp. xviii+456. [DOI](#).
- [6] G. Shimura. “On modular correspondences for $Sp(n, Z)$ and their congruence relations”. *Proc. Nat. Acad. Sci. U.S.A.* **49** (1963), pp. 824–828. [DOI](#).
- [7] T. Tamagawa. “On the ζ -functions of a division algebra”. *Ann. of Math. (2)* **77** (1963), pp. 387–405. [DOI](#).
- [8] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 9.2)*. <https://www.sagemath.org>. 2020.
- [9] K. Vankov. “Explicit Hecke series for symplectic group of genus 4”. *J. Théor. Nombres Bordeaux* **23.1** (2011), pp. 279–298.
- [10] C. Voll. “Functional equations for zeta functions of groups and rings”. *Ann. of Math. (2)* **172.2** (2010), pp. 1181–1218. [DOI](#).